

Co-funded by the Prevention of and Fight against Crime Programme of the European Union



NACIONALNI FORENZIČNI LABORATORIJ

Vodovodna 95 1000 Ljubljana SLOVENIJA

T: +386 (0)1 428 44 93 E: <u>nfl@policija.si</u> www.policija.si

ANALYTICAL REPORT

2-Bromoamphetamine (C9H12BrN)

2-bromo-α-methyl-benzeneethanamine

Remark – other NPS detected: micro trace of amphetamine

Sample ID:	1207-15		
Sample description:	powder - white		
Sample type:	RM-reference material		
Comments ¹ :	Chiron AS Lot#15339RESPONSE -purchasing		
Date of entry:	8/31/2015		

Substance identified- structure ² (base form)	Br NH2		
Systematic name:	2-bromo-α-methyl-benzeneethanamine		
Other names:			
Formula (per base form)	C9H12BrN		
M _w (g/mol)	214,11		
Salt form:	HCI		
StdInChIKey	VGJBPBDFJNFFRO-UHFFFAOYSA-N		
Compound Class	Phenethylamines		
Other NPS detected	micro trace of amphetamine		
Add.info (purity)	98,60%		

² Created by OPSIN free tool: <u>http://opsin.ch.cam.ac.uk/</u> **DOI:** 10.1021/ci100384d

¹ This report has been produced with the financial support of the Prevention of and fight against crime Programme of the European Union (grant agreement number JUST/2013/ISEC/DRUGS/AG/6413). The contents of this report are the sole responsibility of the National Forensic Laboratory and can in no way be taken to reflect the views of the European Commission.





Supporting information


Analytical technique:	applied	remarks
GC-MS (El ionization)	+	NFL GC-RT (min): 3,16
		BP(1): 44; BP(2): 89,BP(3) :90,
FTIR-ATR	+	
FTIR (condensed phase) always		
for the base form of cpd.		

GC-MS (Agilent): GC-method is RT locked to tetracosane (RT=9.53 min). Injection volume 1 ml and split mode (1:50)for GC-MS instruments and 1:5 for GC-MS-FTIR(condensed phase). Injector temperature: 280 $^{\circ}$ C. Column: HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickens 0.25 mm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 $^{\circ}$ C for 1 min, followed by heating up to 293 $^{\circ}$ C at a rate of 18 $^{\circ}$ C/min, hold for 6.1 min, than heating at 50 $^{\circ}$ C/min up to 325 $^{\circ}$ C and finally 2.8 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235 $^{\circ}$ C, source and quadropole temperatures 280 $^{\circ}$ C and 180 $^{\circ}$ C. m/z scan range: from 50 (40) to 550 amu.


FTIR-ATR (Perkin Elmer): scan range 4000-400 cm-1; resolution 4cm-1

FTIR (Spectra analyses-Danny): scan range 4000 to 700, resolution 4cm-1

Abundance



MS-spectrum (EI)



FTIR-ATR

