




ANALYTICAL REPORT

4-HO-McPT (C₁₄H₁₈N₂O)

3-{2-[cyclopropyl(methyl)amino]ethyl}-1H-indol-4-ol

Remark – other NPS detected: **none**

Sample ID:	1961-18
Sample description:	powder
Sample type:	test purchase /ISF projekt (NFL-SI)
Date of sample receipt (DD/MM/YYYY):	21/08/2018
Date of entry (DD/MM/YYYY) into NFL database:	10/09/2018
Report updates (if any) will be published here:	http://www.policija.si/apps/nfl_response_web/seznam.php

Substance identified - structure ¹ (base form)	
Systematic name	3-{2-[cyclopropyl(methyl)amino]ethyl}-1H-indol-4-ol
Other names	
Formula (per base form)	C ₁₄ H ₁₈ N ₂ O
M _w (g/mol)	230,31
Salt form/anions detected	fumarate
StdInChIKey (per base form)	GFVJBFIXZYLPO-UHFFFAOYSA-N
Other NPS detected	none
Additional info (purity..)	> 93 % based on H1 NMR

¹ Created by OPSIN free tool: <http://opsin.ch.cam.ac.uk/> DOI: 10.1021/ci100384d

Report updates

date	comments (explanation)

Instrumental methods (if applied) in NFL

1. GC-MS (Agilent): GC-method is RT locked to tetracosane (9.258 min). Injection volume 1 ml and split mode (1:50). Injector temperature: 280 °C. Chromatographic separation: on column HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickness 0.25 µm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 °C for 1 min, followed by heating up to 190 °C at rate 8 °C/min, then heating up to 293 °C at a rate of 18 °C/min, hold for 7.1 min, then heating at 50 °C/min up to 325 °C and finally 6.1 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235°C, source and quadropole temperatures 280°C and 180°C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300 until 6 min) amu.

2. HPLC-TOF (Agilent): 6230B TOF with Agilent 1260 Infinity HPLC with binary pump, column: Zorbax Eclipse XDB-C18, 50 x 4.6 mm, 1.8 micron. Mobile phases (A) 0.1% formic acid and 1mM ammonium formate in water; (B) 0.1% formic acid in methanol (B). Gradient: starting at 5% B, changing to 40% B over 4 min, then to 70% over 2 min and in 5 min to 100%, hold 1 min and back to 5%, equilibration for 1.7 min. The flow rate: 1.0 ml/min; Injection volume 1 µl. MS parameters: 2GHz, Extended Dynamic range mode to a maximum of 1700 amu, acquisition rate 1.30 spectra/sec. Sample ionisation: by Agilent Jet Stream technology (Dual AJS ESI). Ion source: positive ion scan mode with mass scanning from 82 to 1000 amu. Other TOF parameters: drying gas (N₂) and sheath temperature 325 °C; drying gas flow rate 6 l/min; sheath gas flow rate 8 l/min; nebulizer 25 psig; Vcap. 4000 V; nozzle 2000 V; skimmer 65 V; fragmentor 175 V and Octopole RF 750 V.

3. FTIR-ATR (Perkin Elmer): scan range 4000-400 cm⁻¹; resolution 4cm⁻¹

4. GC- (MS)-IR condensed phase (GC-MS (Agilent) & IR (Spectra analyses-Danny)

GC-method: Injection volume 1 ml and split mode (1:5). Injector temperature 280 °C. Chromatographic separation as above **(1)**. Split MS : IR = 1: 9.

MSD source EI = 70 eV. GC-MS transfer line T= 235°C, source and quadropole temperatures 280°C and 180°C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300) amu.

IR (condensed (solid) phase): IR scan range 4000 to 650, resolution 4 cm⁻¹.

5. IC (anions) (Thermo Scientific, Dionex ICS 2100), Column: IonPac AS19, 2 x 250mm; Eluent: 10mM KOH from 0 to 10 min, 10-58 mM from 10 to 40min; Flow rate: 0.25 ml/min; Temperature: 30°C; Suppressor: AERS 500 2mm, suppressor current 13mA; Inj. Volume: 25 µl


Supporting information

Solubility in	result/remark
CH ₂ Cl ₂	partially
MeOH	partially
H ₂ O	/


Analytical technique:	applied	remarks
GC-MS (EI ionization)	+	NFL GC-RT (min): 7,98 BP(1): 84; BP(2): 159,BP(3) :230,
HPLC-TOF	+	Exact mass (theoretical): 230,1419; measured value Δppm:-0,3; formula:C14H18N2O
FTIR-ATR	+	direct measurement (sample as received)
FTIR (solid phase) always as base form	+	
IC (anions)	+	
NMR (in FKKT)	+	
validation		
other		

ANALYTICAL RESULTS


GC-MS chromatogram (non derivatized sample)




GC-MS chromatogram – TMS derivative




MS (EI)



FTIR-ATR - direct measurement (sample as received fumarate salt)



IR (solid phase – after chromatographic separation)



TOF REPORT


Data File	1961-18.d	Sample Name	ID 1961-18
Sample Type	Sample	Position	P1-C7
Instrument Name	6230B TOF LC-MS	User Name	TG
Acq Method	general-04_12_2017-XDB-C18-ESI+.m	Acquired Time	8/22/2018 4:23:52 PM
IRM Calibration Status	Success	DA Method	a-Drugs_NFL.m
Comment	MeOH		

Compound Table

Label	MFG Formula	Obs. RT	Obs. Mass
Cpd 1: C14 H18 N2 O	C14 H18 N2 O	3.6	230.142

Obs. m/z	Obs. RT	Obs. Mass
231.1492	3.6	230.142

MFE MS Zoomed Spectrum



MS Spectrum Peak List


Obs. m/z	Charge	Abund	Formula	Ion/Isotope
231.1492	1	2563623.5	C14 H18 N2 O	(M+H)+
232.1526	1	370515.27	C14 H18 N2 O	(M+H)+
233.155	1	32647	C14 H18 N2 O	(M+H)+
234.1577	1	2186.53	C14 H18 N2 O	(M+H)+
253.1301	1	1010.84	C14 H18 N2 O	(M+Na)+

--- End Of Report ---

Peak Integration Report


Sample Name:	1961_18_IC	Inj. Vol.:	25,00
Injection Type:	Unknown	Dilution Factor:	1,0000
Program:	ANIONI - julij 2018	Operator:	kemija
Inj. Date / Time:	23-avg-2018 / 09:00	Run Time:	47,10

No.	Time min	Peak Name	Peak Type	Area $\mu\text{S}\cdot\text{min}$	Height μS	Amount mg/L
1,00	33,15	Fumarate	BMB	4,49	11,62	n.a.
		TOTAL:		4,49	11,62	0,00






R E P O R T

Contract No.	C1714-17-460078 (Republic of Slovenia, Ministry of the Interior, POLICE)
Sample ID:	1961-18
Received date:	August 29, 2018
Our notebook code:	NFL-1961-18
NMR sample preparation:	22.2 mg dissolved in 0.7 mL DMSO- d_6
NMR experiments:	^1H , ^{13}C , ^1H - ^1H <i>gs</i> -COSY, ^1H - ^{13}C <i>gs</i> -HSQC, ^1H - ^{13}C <i>gs</i> -HMBC, ^1H - ^{15}N <i>gs</i> -HMBC
Proposed structure with formula, exact mass, molecular weight:	 Chemical Formula: $\text{C}_{14}\text{H}_{18}\text{N}_2\text{O}$ Exact Mass: 230,1419 Molecular Weight: 230,3110
Chemical name:	3-(2-(cyclopropyl(methyl)amino)ethyl)-1 <i>H</i> -indol-4-ol
Comments:	- Structure elucidation based on 1D and 2D NMR spectra and HRMS. - The sample consists of herein identified compound and fumarate in molar ratio of 1 : 0.5, based on ^1H NMR spectrum. - >93% purity of a sample based on ^1H NMR spectrum
Supporting information:	Copies of ^1H and ^{13}C NMR spectra, ^1H and ^{13}C FIDs.
Principal investigator:	Prof. Dr. Janez Košmrlj
Date of report:	September 7, 2018

NFL-1961-18
1H



```


Current Data Parameters
NAME      NFL-1961-18
EXPNO     10
PROCNO    1

F2 - Acquisition Parameters
Date_     20180830
Time      7.26
INSTRUM   spect
PROBHD    5 mm PABBO BB-
PULPROG   zg30
TD         65536
SOLVENT   DMSO
NS         16
DS         2
SWH        10000.000 Hz
FIDRES     0.152588 Hz
AQ         3.2767999 sec
RG         64
DW         50.000 usec
DE         6.50 usec
TE         296.0 K
D1         1.00000000 sec
TD0        1

===== CHANNEL f1 =====
SFO1      500.130885 MHz
NUC1      1H
P1        8.70 usec
PLM1      26.00000000 W

F2 - Processing parameters
SI        65536
SF        500.130041 MHz
WDW       EM
SSB       0
LB        0.30 Hz
GB        0
PC        1.00
  
```

NFL-1961-18
13C



Current Data Parameters
NAME NFL-1961-18
EXPNO 3
PROCNO 1
F2 - Acquisition Parameters
Date_ 20180829
Time 23.07
INSTRUM spect
PROBHD 5 mm PABBO BB-
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 3072
DS 4
SWH 29761.904 Hz
FIDRES 0.454131 Hz
AQ 1.1010048 sec
RG 2050
DM 16.800 usec
DE 6.50 usec
TE 296.0 K
D1 1.0000000 sec
D11 0.0300000 sec
TD0 1
===== CHANNEL f1 =====
SFO1 125.7703637 MHz
NUC1 13C
P1 8.70 usec
PLW1 122.0000000 W
===== CHANNEL f2 =====
SFO2 500.1320005 MHz
NUC2 1H
CDEPRG12 Waltz16
PCPD2 80.00 usec
PLW2 26.00000000 W
PLW12 0.30046001 W
PLW13 0.15113001 W
F2 - Processing parameters
SI 32768
SF 125.7578424 MHz
WDW EM
SSB 0
TB 1.00 Hz
GB 0
PC 1.40