




ANALYTICAL REPORT

EFLEA (C12H17NO3)

N-[1-(2,3-dihydro-1,4-benzodioxin-6-yl)propan-2-yl]-N-methylhydroxylamine

Remark – other NPS detected:

Sample ID:	1866-17
Sample description:	powder
Sample type:	test purchase /ISF projekt (NFL-SI)
Date of sample receipt (M/D/Y):	10/18/2017
Date of entry (M/D/Y) into NFL database:	1/9/2018
Report updates (if any) will be published here:	http://www.policija.si/apps/nfl_response_web/seznam.php

Substance identified - structure ¹ (base form)	
Systematic name	N-[1-(2,3-dihydro-1,4-benzodioxin-6-yl)propan-2-yl]-N-methylhydroxylamine
Other names	N-hydroxy-3,4-ethylenedioxy-N-methylamphetamine; N-Hydroxy-N-methyl-1-(2,3-dihydro-1,4-benzodioxin-6-yl)propane-2-amine; 1-(2,3-Dihydro-1,4-benzodioxin-6-yl)-N-hydroxy-N-methylpropan-2-amine
Formula (per base form)	C12H17NO3
M _w (g/mol)	223,27
Salt form/anions detected	HCl
StdInChIKey (per base form)	FFWSMXIWPUUWNK-UHFFFAOYSA-N
Other NPS detected	
Additional info (purity..)	pure by HPLC-TOF, NMR

¹ Created by OPSIN free tool: <http://opsin.ch.cam.ac.uk/> DOI: 10.1021/ci100384d

Report updates

date	comments (explanation)

Instrumental methods (if applied) in NFL

1. GC-MS (Agilent): GC-method is RT locked to tetracosane (9.258 min). Injection volume 1 ml and split mode (1:50). Injector temperature: 280 °C. Chromatographic separation: on column HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickness 0.25 µm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 °C for 1 min, followed by heating up to 190 °C at rate 8 °C/min, then heating up to 293 °C at a rate of 18 °C/min, hold for 7.1 min, then heating at 50 °C/min up to 325 °C and finally 6.1 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235 °C, source and quadropole temperatures 280 °C and 180 °C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300 until 6 min) amu.

2. HPLC-TOF (Agilent): 6230B TOF with Agilent 1260 Infinity HPLC with binary pump, column: Zorbax Eclipse XDB-C18, 50 x 4.6 mm, 1.8 micron. Mobile phases (A) 0.1% formic acid and 1mM ammonium formate in water; (B) 0.1% formic acid in methanol (B). Gradient: starting at 5% B, changing to 40% B over 4 min, then to 70% over 2 min and in 5 min to 100%, hold 1 min and back to 5%, equilibration for 1.7 min. The flow rate: 1.0 ml/min; Injection volume 1 µl. MS parameters: 2GHz, Extended Dynamic range mode to a maximum of 1700 amu, acquisition rate 1.30 spectra/sec. Sample ionisation: by Agilent Jet Stream technology (Dual AJS ESI). Ion source: positive ion scan mode with mass scanning from 82 to 1000 amu. Other TOF parameters: drying gas (N₂) and sheath temperature 325 °C; drying gas flow rate 6 l/min; sheath gas flow rate 8 l/min; nebulizer 25 psig; Vcap. 4000 V; nozzle 2000 V; skimmer 65 V; fragmentor 175 V and Octopole RF 750 V.

3. FTIR-ATR (Perkin Elmer): scan range 4000-400 cm⁻¹; resolution 4cm⁻¹

4. GC- (MS)-IR condensed phase (GC-MS (Agilent) & IR (Spectra analyses-Danny)

GC-method: Injection volume 1 ml and split mode (1:5). Injector temperature 280 °C. Chromatographic separation as above **(1)**. Split MS : IR = 1: 9.

MSD source EI = 70 eV. GC-MS transfer line T= 235 °C, source and quadropole temperatures 280 °C and 180 °C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300) amu.

IR (condensed (solid) phase): IR scan range 4000 to 650, resolution 4 cm⁻¹.

5. IC (anions) (Thermo Scientific, Dionex ICS 2100), Column: IonPac AS19, 2 x 250mm; Eluent: 10mM from 0 to 10 min, 10-58 mM from 10 to 40min; Flow rate: 0.25 ml/min; Temperature: 30 °C; Suppressor: AERS 500 2mm, suppressor current 13mA; Inj. Volume: 25 µl

Supporting information



Solubility in	result/remark
CH ₂ Cl ₂	partially
MeOH	soluble
H ₂ O	partially

Analytical technique:	applied	remarks
GC-MS (EI ionization)	+	NFL GC-RT (min): 6,76 (as TMS derivative) BP(1): 146; BP(2): 149,BP(3) :147,
HPLC-TOF	+	Exact mass (theoretical): 223,1208; measured value Δppm:1,07; formula:C12H17NO3
FTIR-ATR	+	direct measurement (sample as received)
FTIR (condensed phase) always as base form	-	
IC (anions)	+	
NMR (in FKKT)	+	
validation		
other		


ANALYTICAL RESULTS

MS (EI) of EFLEA-TMS derivative

Abundance



FTIR-ATR - direct measurement (sample as received)



TOF REPORT


Data File	1866-17.d	Sample Name	1866-17
Sample Type	Sample	Position	P2-A5
Instrument Name	6230B TOF LC-MS	User Name	TG
Acq Method	general-19_07_2017-XDB-C18-ESI-final.m	Acquired Time	10/20/2017 9:18:13 AM
IRM Calibration Status	Success	DA Method	Drugs_NFL.m
Comment	MeOH		

Compound Table


Label	Compound Name	MFG Formula	Obs. RT	Obs. Mass
Cpd 2: EFLEA	EFLEA	C12 H17 N O3	5.18	223.1206

Name	Obs. m/z	Obs. RT	Obs. Mass	DB RT	DB Formula	DB Mass	DB Mass Error (ppm)
EFLEA	224.1281	5.18	223.1206	5.18	C12 H17 N O3	223.1208	1.07


Compound Chromatograms



MFE MS Zoomed Spectrum



MS Zoomed Spectrum



MS Spectrum Peak List


Obs. m/z	Charge	Abund	Formula	Ion/Isotope
224.1281	1	13944696	C12 H17 N O3	(M+H)+
225.1298	1	1776243.09	C12 H17 N O3	(M+H)+
226.1321	1	179255.35	C12 H17 N O3	(M+H)+
227.1346	1	15704.16	C12 H17 N O3	(M+H)+
262.0832	1	4273.74		(M+K)+
263.0771	1	1348.45		(M+K)+
447.2468	1	102871.11	C12 H17 N O3	(2M+H)+
448.2503	1	26225.25	C12 H17 N O3	(2M+H)+
449.2529	1	4849.88	C12 H17 N O3	(2M+H)+

--- End Of Report ---

Peak Integration Report

Sample Name:	1866-17	Inj. Vol.:	25,00
Injection Type:	Unknown	Dilution Factor:	1,0000
Program:	ANIONI	Operator:	kemija
Inj. Date / Time:	20-okt-2017 / 11:25	Run Time:	42,00

No.	Time min	Peak Name	Peak Type	Area $\mu\text{S}\cdot\text{min}$	Height μS	Amount mg/L
1,00	9,35	Chloride	BMB	3,17	14,72	n.a.
		TOTAL:		3,17	14,72	0,00



University
of Ljubljana
Faculty of Chemistry
and Chemical Technology




R E P O R T

Contract No.	C1714-17-460078 (Republic of Slovenia, Ministry of the Interior, POLICE)
Sample ID:	1866-17
Received date:	December 8, 2017
Our notebook code:	NFL-1866-17
NMR sample preparation:	16.5 mg dissolved in 0.7 mL DMSO- <i>d</i> ₆
NMR experiments:	¹ H, ¹³ C, ¹ H- ¹ H <i>gs</i> -COSY, ¹ H- ¹³ C <i>gs</i> -HSQC, ¹ H- ¹³ C <i>gs</i> -HMBC, ¹ H- ¹⁵ N <i>gs</i> -HMBC
Proposed structure, formula, exact mass, molecular weight:	 Chemical Formula: C ₁₂ H ₁₈ NO ₃ ⁺ Exact Mass: 224,1281 Molecular Weight: 224,2795
Chemical name:	N-protonated N-(1-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)propan-2-yl)-N-methylhydroxylamine
Comments:	- Structure elucidation based on 1D and 2D NMR spectra and HRMS. - >98% purity of the sample based on ¹H NMR spectrum
Supporting information:	Copies of ¹ H and ¹³ C NMR spectra, ¹ H and ¹³ C FIDs
Principal investigator:	Prof. Dr. Janez Košmrlj
Date of report:	December 22, 2017

NFL-1866-17
1H

13.196
11.529




Current Data Parameters
NAME NFL-1866-17
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20171211
Time 17.55
INSTRUM spect
PROBHD 5 mm PABBO BB-
PULPROG zg30
TD 65536
SOLVENT DMSO
NS 16
DS 2
SWH 10000.000 Hz
FIDRES 0.152588 Hz
AQ 3.2767999 sec
RG 80.6
DW 50.000 usec
DE 6.50 usec
TE 296.0 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
SFO1 500.130885 MHz
NUC1 1H
P1 8.70 usec
PLW1 26.00000000 W
F2 - Processing parameters
SI 65536
SF 500.1300005 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

NFL-1866-17 13C



Current Data Parameters
NAME NFL-1866-17
EXPNO 8
PROCNO 1

F2 - Acquisition Parameters
Date_ 20171215
Time 7.31
INSTRUM spect
PROBHD 5 mm PABBO BB-
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 5120
DS 4
SWH 29761.904 Hz
FIDRES 0.454131 Hz
AQ 1.1010048 sec
RG 2050
DW 16.800 usec
DE 6.50 usec
TE 296.0 K
D1 2.00000000 sec
D11 0.03000000 sec
TD0 1

===== CHANNEL f1 =====
SF01 125.7703637 MHz
NUC1 13C
P1 8.70 usec
PLM1 122.0000000 W

===== CHANNEL f2 =====
SF02 500.1320005 MHz
NUC2 1H
CPDPRG2 waltz16
PCPD2 80.00 usec
PLM2 26.00000000 W
PLM12 0.30046001 W
PLM13 0.15113001 W

F2 - Processing parameters
SI 32768
SF 125.7578438 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40