




## ANALYTICAL REPORT

### NM-2201 (C<sub>24</sub>H<sub>22</sub>FNO<sub>2</sub>)

#### naphthalen-1-yl 1-(5-fluoropentyl)-1H-indole-3-carboxylate

Remark – other NPS detected: **none**

Sample ID:	1776-17
Sample description:	powder - white off
Sample type:	seized /NM
Date of sample receipt (M/D/Y):	1/31/2017
Date of entry (M/D/Y) into NFL database:	2/23/2017
Report <sup>1</sup> updates (if any) will be published here:	<a href="http://www.policija.si/apps/nfl_response_web/seznam.php">http://www.policija.si/apps/nfl_response_web/seznam.php</a>

Substance identified - structure <sup>2</sup> (base form)	
Systematic name	naphthalen-1-yl 1-(5-fluoropentyl)-1H-indole-3-carboxylate
Other names	CBL-2201, 5-Fluoro-SDB-005 Indole
Formula (per base form)	C <sub>24</sub> H <sub>22</sub> FNO <sub>2</sub>
M <sub>w</sub> (g/mol)	375,44
Salt form/anions detected	base
StdInChIKey (for base form)	PRGFSQYZCKBQO-UHFFFAOYSA-N
Other NPS detected	none
Additional info (purity..)	Sample is pure by GC-MS, TOF

<sup>1</sup> This report has been produced with the financial support of the Prevention of and fight against crime Programme of the European Union (grant agreement number JUST/2013/ISEC/DRUGS/AG/6413). The contents of this report are the sole responsibility of the National Forensic Laboratory and can in no way be taken to reflect the views of the European Commission.

<sup>2</sup> Created by OPSIN free tool: <http://opsin.ch.cam.ac.uk/> DOI: 10.1021/ci100384d

## Report updates

date	comments (explanation)

### Instrumental methods (if applied) in NFL

**1. GC-MS** (Agilent): GC-method is RT locked to tetracosane (9.258 min). Injection volume 1 ml and split mode (1:50). Injector temperature: 280 °C. Chromatographic separation: on column HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickness 0.25 µm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 °C for 1 min, followed by heating up to 190 °C at rate 8 °C/min, then heating up to 293 °C at a rate of 18 °C/min, hold for 7.1 min, then heating at 50 °C/min up to 325 °C and finally 6.1 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235 °C, source and quadropole temperatures 280 °C and 180 °C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300 until 6 min) amu.

**2. HPLC-TOF** (Agilent): 6230B TOF with Agilent 1260 Infinity HPLC with binary pump, column: Zorbax Eclipse XDB-C18, 50 x 4.6 mm, 1.8 micron. Mobile phases (A) 0.1% formic acid and 1mM ammonium formate in water; (B) 0.1% formic acid in methanol (B). Gradient: starting at 5% B, changing to 40% B over 4 min, then to 70% over 2 min and in 5 min to 100%, hold 1 min and back to 5%, equilibration for 1.7 min. The flow rate: 1.0 ml/min; Injection volume 1 µl. MS parameters: 2GHz, Extended Dynamic range mode to a maximum of 1700 amu, acquisition rate 1.30 spectra/sec. Sample ionisation: by Agilent Jet Stream technology (Dual AJS ESI). Ion source: positive ion scan mode with mass scanning from 82 to 1000 amu. Other TOF parameters: drying gas (N<sub>2</sub>) and sheath temperature 325 °C; drying gas flow rate 6 l/min; sheath gas flow rate 8 l/min; nebulizer 25 psig; Vcap. 4000 V; nozzle 2000 V; skimmer 65 V; fragmentor 175 V and Octopole RF 750 V.

**3. FTIR-ATR** (Perkin Elmer): scan range 4000-400 cm<sup>-1</sup>; resolution 4cm<sup>-1</sup>

**4. GC- (MS)-IR** condensed phase (GC-MS (Agilent) & IR (Spectra analyses-Danny)

GC-method: Injection volume 1 ml and split mode (1:5). Injector temperature 280 °C. Chromatographic separation as above (**1**). Split MS : IR = 1: 9.

MSD source EI = 70 eV. GC-MS transfer line T= 235 °C, source and quadropole temperatures 280 °C and 180 °C, respectively. Scan range m/z scan range: from 50 (30 until 6 min.) to 550 (300) amu.

IR (condensed (solid) phase): IR scan range 4000 to 650, resolution 4 cm<sup>-1</sup>.

**5. IC** (anions) (Thermo Scientific, Dionex ICS 2100), Column: IonPac AS19, 2 x 250mm; Eluent: 10mM from 0 to 10 min, 10-58 mM from 10 to 40min; Flow rate: 0.25 ml/min; Temperature: 30 °C; Suppressor: AERS 500 2mm, suppressor current 13mA; Inj. Volume: 25 µl

## Supporting information


Solubility in	result/remark
CH <sub>2</sub> Cl <sub>2</sub>	soluble
MeOH	partially
H <sub>2</sub> O	partially

Analytical technique:	applied	remarks
GC-MS (EI ionization)	+	NFL GC-RT (min): 17,08 BP(1): 232; BP(2): 144,BP(3) :212,
HPLC-TOF	+	Exact mass (theoretical): 375,1635; measured value Δppm:-0,38; formula:C <sub>24</sub> H <sub>22</sub> FNO <sub>2</sub>
FTIR-ATR	+	direct measurement (sample as received)
FTIR (condensed phase) always as base form	+	
IC (anions)	+	
NMR (in FKKT)	-	
validation	+	MS consistent by spectra published in SWGDRUG.L, ENFSI 15.L, DD2016.L and Cayman.L; FTIR ATR spectrum in line with the published SWGDURG library (correlation > 0.997)
other		


# ANALYTICAL RESULTS

MS (EI)


Abundance



FTIR-ATR - direct measurement (sample as received)



IR (condensed phase – after chromatographic separation)



# TOF REPORT


<b>Data File</b>	NM-2201_1776-17.d	<b>Sample Name</b>	1776-17
<b>Sample Type</b>	Sample	<b>Position</b>	P1-A2
<b>Instrument Name</b>	6230B TOF LC-MS	<b>User Name</b>	TG
<b>Acq Method</b>	general-20_12_2016-XDB-C18-ESI-poz-soft.m	<b>Acquired Time</b>	2/22/2017 1:11:34 PM
<b>IRM Calibration Status</b>	Success	<b>DA Method</b>	Drugs_NFL.m
<b>Comment</b>	MeOH		

**Compound Table**


Label	Compound Name	MFG Formula	Obs. RT	Obs. Mass
Cpd 9: NM-2201	NM-2201	C24 H22 F N O2	10.16	375.1636

Name	Obs. m/z	Obs. RT	Obs. Mass	DB RT	DB Formula	DB Mass	DB Mass Error (ppm)
NM-2201	376.1708	10.16	375.1636	10.16	C24 H22 F N O2	375.1635	-0.38


**Compound Chromatograms**



**MFE MS Zoomed Spectrum**



**MS Zoomed Spectrum**



**MS Spectrum Peak List**

Obs. m/z	Charge	Abund	Formula	Ion/Isotope
376.1708	1	4150424	C24 H22 F N O2	(M+H)+
377.1746	1	1124551.69	C24 H22 F N O2	(M+H)+
393.1975	1	361656.5	C24 H22 F N O2	(M+NH4)+
398.153	1	494640.06	C24 H22 F N O2	(M+Na)+
768.3612	1	2488874.5		(2M+NH4)+
769.3647	1	1364877.08		(2M+NH4)+
770.368	1	355182.8		(2M+NH4)+
773.3168	1	1638165.13		(2M+Na)+
774.3206	1	863383.92		(2M+Na)+
789.2911	1	588899.56		(2M+K)+

--- End Of Report ---

### Peak Integration Report

Sample Name:	1776-17_IC	Inj. Vol.:	25,00
Injection Type:	Unknown	Dilution Factor:	1,0000
Program:	ANIONI	Operator:	kemija
Inj. Date / Time:	22-feb-2017 / 11:03	Run Time:	42,00

No.	Time min	Peak Name	Peak Type	Area $\mu\text{S}\cdot\text{min}$	Height $\mu\text{S}$	Amount n.a.
		TOTAL:		0,00	0,00	0,00

