



ANALYTICAL REPORT¹

SDB-005 (
C23H22N2O2)

naphthalen-1-yl 1-pentyl-1H-indazole-3-carboxylate

Remark – other NPS detected: **none**

Sample ID:	1389-15
Sample description:	powder - brownish-light
Sample type:	test purchase /RESPONSE -purchasing
Date of sample receipt (M/D/Y):	12/9/2015
Date of entry (M/D/Y) into NFL database:	12/18/2015
Report updates (if any) will be published here:	http://www.policija.si/apps/nfl_response_web/seznam.php

Substance identified - structure ² (base form)	
Systematic name	naphthalen-1-yl 1-pentyl-1H-indazole-3-carboxylate
Other names	
Formula (per base form)	C23H22N2O2
M _w (g/mol)	358,44
Salt form/anions detected	base
StdInChIKey	JBVNFKZVDLFOAY-UHFFFAOYSA-N
Compound Class	Cannabinoids
Other NPS detected	none
Add.info (purity..)	pure by GC, HPLC-TOF

¹ This report has been produced with the financial support of the Prevention of and fight against crime Programme of the European Union (grant agreement number JUST/2013/ISEC/DRUGS/AG/6413). The contents of this report are the sole responsibility of the National Forensic Laboratory and can in no way be taken to reflect the views of the European Commission.

² Created by OPSIN free tool: <http://opsin.ch.cam.ac.uk/> DOI: 10.1021/ci100384d

Report updates

date	comments (explanation)

Instrumental methods (if applied) in NFL

1. GC-MS (Agilent): GC-method is RT locked to tetracosane (RT=9.53 min). Injection volume 1 ml and split mode (1:50) . Injector temperature: 280 °C. Chromatographic separation: on column HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickness 0.25 mm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 °C for 1 min, followed by heating up to 293 °C at a rate of 18 °C/min, hold for 6.1 min, then heating at 50 °C/min up to 325 °C and finally 6.1 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235°C, source and quadropole temperatures 280°C and 180°C, respectively. Scan range m/z scan range: from 50 (40) to 550 amu.

2. HPLC-TOF (Agilent): 6230B TOF with Agilent 1260 Infinity HPLC with binary pump, column: Zorbax Eclipse XDB-C18, 50 x 4.6 mm, 1.8 micron. Mobile phases (A) 0.1% formic acid and 1mM ammonium formate in water; (B) 0.1% formic acid in methanol (B). Gradient: starting at 5% B, changing to 40% B over 4 min, then to 70% over 2 min and in 5 min to 100%, hold 1 min and back to 5%, equilibration for 1.7 min. The flow rate: 1.0 ml/min; Injection volume 1 µl. MS parameters: 2GHz, Extended Dynamic range mode to a maximum of 1700 amu, acquisition rate 1.30 spectra/sec. Sample ionisation: by Agilent Jet Stream technology (Dual AJS ESI). Ion source: positive ion scan mode with mass scanning from 82 to 1000 amu. Other TOF parameters: drying gas (N2) and sheath temperature 325 °C; drying gas flow rate 6 l/min; sheath gas flow rate 8 l/min; nebulizer 25 psig; Vcap. 4000 V; nozzle 2000 V; skimmer 65 V; fragmentor 175 V and Octopole RF 750 V.

3. FTIR-ATR (Perkin Elmer): scan range 4000-400 cm⁻¹; resolution 4cm⁻¹

4. GC- (MS)-IR condensed phase (GC-MS (Agilent) & IR (Spectra analyses-Danny)

GC-method: Injection volume 1 ml and split mode (1:5). Injector temperature 280 °C. Chromatographic separation as above (**1**). Split MS : IR = 1:9.

MSD source EI = 70 eV. GC-MS transfer line T= 235°C, source and quadropole temperatures 280°C and 180°C, respectively. Scan range m/z scan range: from 50 (40) to 550 amu.

IR (condensed phase): IR scan range 4000 to 650, resolution 4 cm⁻¹.

5. IC (anions) (Thermo Scientific, Dionex ICS 2100), Column: IonPac AS19, 2 x 250mm; Eluent: 10mM from 0 to 10 min, 10-58 mM from 10 to 40min; Flow rate: 0.25 ml/min; Temperature: 30°C; Suppressor: AERS 500 2mm, suppressor current 13mA; Inj. Volume: 25 µl

Supporting information


Solubility in	result/remark
CH ₂ Cl ₂	soluble
MeOH	low (bad)
H ₂ O	low (bad)

Analytical technique:	applied	remarks
GC-MS (EI ionization)	+	NFL GC-RT (min): 14,61 BP(1): 215; BP(2): 145,BP(3) :216,
HPLC-TOF	+	Exact mass (theoretical): 358,1681; measured value Δppm:-0,16; formula: C ₂₃ H ₂₂ N ₂ O ₂
FTIR-ATR	+	direct measurement (sample as received)
FTIR (condensed phase) always as base form	+	
IC (anions)	+	
NMR (in FKKT)	+	
validation		confirmed by reference material
other		


ANALYTICAL RESULTS

MS (EI)


Abundance



FTIR-ATR - direct measurement (sample as received)



IR (condensed phase – after chromatographic separation)



TOF REPORT


Data File	SDB005_1389-15_TOF.d	Sample Name	ID_1389-15
Sample Type	Sample	Position	P1-A9
Instrument Name	6230B TOF LC-MS	User Name	TG
Acq Method	general-17112015-XDB-C18-ESI-poz.m	Acquired Time	12/11/2015 12:09:32 PM
IRM Calibration Status	Success	DA Method	Drugs_NFL.m
Comment	extract in MeOH		

Compound Table


Label	Compound Name	MFG Formula	Obs. RT	Obs. Mass
Cpd 9: SDB-005	SDB-005	C23 H22 N2 O2	10.77	358.1682

Name	Obs. m/z	Obs. RT	Obs. Mass	DB RT	DB Formula	DB Mass	DB Mass Error (ppm)
SDB-005	359.1756	10.77	358.1682	10.77	C23 H22 N2 O2	358.1681	-0.16


Compound Chromatograms



MFE MS Zoomed Spectrum



MS Zoomed Spectrum



MS Spectrum Peak List

Obs. m/z	Charge	Abund	Formula	Ion/Isotope
359.1756	1	4609515.5	C23 H22 N2 O2	(M+H)+
360.1794	1	1187359.34	C23 H22 N2 O2	(M+H)+
361.1821	1	151761	C23 H22 N2 O2	(M+H)+
381.1575	1	112617.91	C23 H22 N2 O2	(M+Na)+
717.3434	1	2840886.5	C23 H22 N2 O2	(2M+H)+
718.3471	1	1471777.52	C23 H22 N2 O2	(2M+H)+
719.3504	1	390228.05	C23 H22 N2 O2	(2M+H)+
739.3261	1	905828.56	C23 H22 N2 O2	(2M+Na)+
740.3292	1	459496.43	C23 H22 N2 O2	(2M+Na)+
741.3318	1	111823.14	C23 H22 N2 O2	(2M+Na)+

--- End Of Report ---

Peak Integration Report

Sample Name:	SDB-005_1389-15_IC	Inj. Vol.:	25,00
Injection Type:	Unknown	Dilution Factor:	1,0000
Program:	ANIONI	Operator:	kemija
Inj. Date / Time:	11-dec-2015 / 18:05	Run Time:	42,00

No.	Time min	Peak Name	Peak Type	Area $\mu\text{S}\cdot\text{min}$	Height μS	Amount n.a.
		TOTAL:		0,00	0,00	0,00

