

SLOVENIJA



T: +386 (0)1 428 44 93 E: nfl@policija.si www.policija.si

ANALYTICAL REPORT

para-Methyl-4-methylaminorex (C11H14N2O)

4,5-Dihydro-4-methyl-5-(4-methylphenyl)-2-oxazolamine

Remark – other NPS detected: none

Sample ID:	1240-15		
Sample description:	crystal - white		
Sample type:	RM-reference material		
Comments ¹ :	Chiron AS Lot#14127RESPONSE -purchasing		
Date of entry:	8/31/2015		

Substance identified- structure ² (base form)	N NH2
Systematic name:	4,5-Dihydro-4-methyl-5-(4-methylphenyl)-2-oxazolamine
Other names:	
Formula (per base form)	C11H14N2O
M _w (g/mol)	190,24
Salt form:	base
StdInChIKey	NPILLHMQNMXXTL-UHFFFAOYSA-N
Compound Class	Others
Other NPS detected	none
Add.info (purity)	99,50%

¹ This report has been produced with the financial support of the Prevention of and fight against crime Programme of the European Union (grant agreement number JUST/2013/ISEC/DRUGS/AG/6413). The contents of this report are the sole responsibility of the National Forensic Laboratory and can in no way be taken to reflect the views of the European Commission.

² Created by OPSIN free tool: http://opsin.ch.cam.ac.uk/ **DOI:** 10.1021/ci100384d






Supporting information


Analytical technique:	applied	remarks
GC-MS (El ionization)	+	NFL GC-RT (min): 5,02
		BP(1): 70; BP(2): 43,BP(3) :146,
FTIR-ATR	+	direct measurement
FTIR (condensed phase) always for the base form of cpd.	+	

GC-MS (Agilent): GC-method is RT locked to tetracosane (RT=9.53 min). Injection volume 1 ml and split mode (1:50) for GC-MS instruments and 1:5 for GC-MS-FTIR(condensed phase). Injector temperature: 280 $^{\circ}$ C. Column: HP1-MS (100% dimethylpolysiloxane), length 30 m, internal diameter 0.25 mm, film thickens 0.25 mm. Carrier gas He: flow-rate 1.2 ml/min. GC oven program: 170 $^{\circ}$ C for 1 min, followed by heating up to 293 $^{\circ}$ C at a rate of 18 $^{\circ}$ C/min, hold for 6.1 min, than heating at 50 $^{\circ}$ C/min up to 325 $^{\circ}$ C and finally 2.8 min isothermal. MSD source EI = 70 eV. GC-MS transfer line T= 235 $^{\circ}$ C, source and quadropole temperatures 280 $^{\circ}$ C and 180 $^{\circ}$ C. m/z scan range: from 50 (40) to 550 amu.

FTIR-ATR (Perkin Elmer): scan range 4000-400 cm-1; resolution 4cm-1 FTIR (Spectra analyses-Danny): scan range 4000 to 700, resolution 4cm-1



GC-MS(EI)



FTIR-ATR direct measurement

